

Mini-Review

Apoptosis and in vitro Alzheimer disease neuronal models

P. Calissano,^{1,2} C. Matrone¹ and G. Amadoro^{1,*}

¹Institute of Neurobiology and Molecular Medicine; CNR; ²European Brain Research Institute (EBRI); Rome, Italy

Key words: apoptosis, caspase, amyloid beta, tau protein, neurotrophin, Alzheimer disease

Alzheimer disease (AD) is a human neurodegenerative disease characterized by co-existence of extracellular senile plaques (SP) and neurofibrillary tangles (NFT) associated with an extensive neuronal loss, primarily in the cerebral cortex and hippocampus. Several studies suggest that caspase(s)-mediated neuronal death occurs in cellular and animal AD models as well as in human brains of affected patients, although an etiologic role of apoptosis in such neurodegenerative disorder is still debated. This review summarizes the experimental evidences corroborating the possible involvement of apoptosis in AD pathogenesis and discusses the usefulness of ad hoc devised in vitro approaches to study how caspase(s), amyloidogenic processing and tau metabolism might reciprocally interact leading to neuronal death.

Introduction

Alzheimer disease (AD) is the most common human late-onset and sporadic neurodegenerative disorder characterized by global cognitive decline including a progressive loss of memory, orientation and reasoning. The neuropathological hallmarks of AD include synaptic loss and/or dysfunction, diminished neuronal metabolism, senile plaques (SPs), neurofibrillary tangles (NFTs) and loss of multiple neurotransmitter systems.¹

SPs typically consist of aggregated amyloid beta (A β), abnormal neurites and glial cells. The accumulation of A β due to a dysregulated proteolytic processing of its precursor molecule, the Amyloid Precursor Protein (APP), exerts a crucial role in neuronal loss or dysfunction through a cascade of events which include oxidative stress, membrane damage, altered mitochondrial metabolism, abortive cell cycle events, Ca⁺⁺ imbalance, protein misfolding, DNA damage/repair and inflammatory processes.²

NFTs are intracellular accumulations of cytoskeletal elements, largely made of Paired Helical Filaments (PHF), whose main constituent is abnormally phosphorylated tau. Tangles could potentially damage neurons by disrupting transport of various cellular components, including that of Nerve Growth Factor (NGF)-receptor complex, thus leading to degeneration of the tangle-bearing neurons.³

Thanks to the experimental work carried out in hundreds of laboratories, it has been unequivocally demonstrated that both APP and tau proteins play a crucial role in the onset of AD. Moreover, several strong genetic evidences corroborate the “amyloid cascade hypothesis” according to which A β production is the trigger factor affecting downstream tau metabolism.⁴ Mutations in several known genes linked to AD familial forms (APP, presenilin-1 or presenilin-2 gene) and genetic or environmental risk factors (Apolipoprotein E 4 variant and metals or pesticides exposure) alter A β cellular processing or its properties, leading to an increase of the A β _{42/40} ratio or its propensity to aggregate.¹ Moreover, A β causes caspases-mediated tau cleavage and hyperphosphorylation by activating specific kinases, thus promoting its aggregation, mis-localization and accumulation with consequent NFTs formation.⁵

Although it is still unclear why specific vulnerable neuronal population, with special emphasis to forebrain cholinergic neurons which provide the majority of cholinergic innervations to cerebral cortex and hippocampus, die in the brain of AD patients, a growing number of studies actually indicate apoptosis as possible initial trigger of the pathology.^{6,7}

In this review we will summarize the current findings regarding this hypothesis and we will discuss the convenience of ad hoc devised in vitro models to dissect the single molecular steps linking apoptosis with A β production and tau altered processing. A special emphasis will be devoted to analyze the possible crucial role of NGF and other neurotrophins, since the evidences demonstrating its involvement in the onset of AD are becoming conspicuous.⁸

Alzheimer Disease and Apoptotic Events

Several studies presently indicate that apoptosis might occur in, and contribute to, AD onset and progression.⁷ Stimuli for apoptosis in AD include increased oxidative stress, dysregulation of ion homeostasis, growth factor deprivation, accumulation of A β , metabolic impairment, reduced clearance of toxin, mitochondrial dysfunction, DNA damage, protein aggregation.^{9,7} Nevertheless, while the role of apoptosis in in vitro models and transgenic animal models of neurodegeneration has been largely documented, its occurrence and role in human postmortem AD brain is controversial. Despite a growing number of studies underlying caspases and apoptosis involvement in AD, no direct role of apoptotic death in AD etiology has still been proven although the presence of apoptotic bodies, DNA fragmentation, granulated and marginated chromatin and shrunken and irregular cell shapes have been largely reported in tissue sections of brains from affected patients.^{10,11} Moreover, an imbalanced level

*Correspondence to: G. Amadoro; Institute of Neurobiology and Molecular Medicine; CNR; Via del Fosso di Fiorano 64-65; Rome 00143 Italy; Tel.: +39.06501703269; Fax: +3906501703313; Email: g.amadoro@inmm.cnr.it

Submitted: 12/19/08; Accepted: 12/23/08

Previously published online as a *Communicative & Integrative Biology* E-publication: <http://www.landesbioscience.com/journals/cib/article/7704>

of some molecular apoptotic markers such as pro-apoptotic (Bax, Bak and Bad) and anti-apoptotic (Bcl-2 and Bcl-x_L) proteins—members of Bcl-2 protein family^{12,13}—and the initiator caspases 8 and 9 and the effector caspases 3 and 6 have been also reported in post-mortem brains of AD patients.^{11,14–20} Moreover, expression profiling analysis of thousands of genes in brain tissue samples from AD and age-matched control patients has revealed a marked decrease in expression of some anti-apoptotic gene such as NCKAP1.²¹ In addition, immunohistochemical and biochemical studies report the presence of active caspase(s) and caspase-cleaved substrates in neurons, around senile plaques and neurofibrillary tangles^{10,11,22,23} and also in postsynaptic densities.²⁴ Both caspase-cleaved APP and activated caspase 3 have been shown to be present and associated to granulovacuolar degeneration, a diagnostic AD neuropathological sign in brains of affected patients.²⁵ Finally, a marked co-localization of pathological hyperphosphorylated tau, cleaved caspase-3 and caspase-6 have been recently reported in TUNEL-positive neurons in the brainstem of AD patients.²⁶

Caspases and APP. Caspases have a direct role in amyloid precursor protein (APP) processing and in the biogenesis of A β peptide species.²⁷ Particularly, the C31 C-terminal peptide obtained by caspase-3 mediated APP cleavage seems to mediate apoptosis by transcriptional regulation of some genes.²⁸ Caspase-3 mediated APP cleavage also stabilizes BACE—the β -secretase enzyme initiating the APP cleavage to produce A β peptide—which accumulates in endosomes, where increases A β production.²⁹

Exposure of cultured cortical neurons to A β or infection of rat hippocampal neurons with APP-expressing adenovirus which causes an A β accumulation, induces activation of caspase-3 and apoptosis,^{30–33} suggesting that caspase(s) not only participate in the generation of A β but they may also directly mediate its toxic effect on neuronal survival.³⁴

As will be discussed below, APP-derived toxic peptides may not only originate by apoptosis activation but may also be responsible of it in viable neurons. Thus, APP-derived A β peptides can activate caspases through the extrinsic pathway, implicating binding of extracellular A β to cell sites, while other studies suggest that the intrinsic pathway may be more relevant.³⁵ Intracellular accumulation of A β in endoplasmic reticulum or endosomes may activate apoptotic mechanism(s) through the unfolded protein response (URP) or endoplasmic reticulum stress.³⁶ Alternatively, intracellular A β may bind to alcohol dehydrogenase within mitochondria and activates apoptosis causing mitochondrial stress.³⁷ Interaction of A β with mitochondrial Cyclophilin D causes synaptic damage observed in AD and absence of Cyclophilin D protects neurons from A β - and oxidative stress-induced apoptotic cell death.³⁸ A β 1–42 also impairs proteasome activity and A β immunotherapy rescues the proteasome dysfunction reported in 3X transgenic AD animal models thus confirming that its intracellular accumulation alters the ubiquitin-proteasomal system *in vivo*.³⁹ A β upregulates the intracellular levels of E2-25K/Hip-2, an E2 ubiquitin-conjugating enzyme, which stabilizes endoplasmic reticulum (ER)-resident caspase-12 protein by inhibiting proteasome activity.⁴⁰

Pharmacological or molecular inhibition of particular members of the caspases family, such as caspase 2, 3, 8 and 12 has been reported to offer partial or complete protection against A β -induced apoptotic cell death *in vitro*.^{41–45}

As far as the effect of caspase(s) inhibition on APP metabolism in cellular and animal models, it has been reported that specific down-regulation of caspase-6 in human primary neuronal cultures prevents serum-deprivation mediated A β increase, as well as *in vitro* cell death.⁴⁶ In a similar fashion, IETD, a caspase III inhibitor including caspase-6, -8 and -9 prevents APP cleavage in staurosporine-induced cell death in COS transfected cells.⁴⁷ In agreement with previous *in vitro* experimental data, caspase inhibition *in vivo* by bafilomicin, a pan-caspase(s) inhibitor, abolishes brain trauma-induced increase in A β and reduces neuronal degeneration in hippocampus of injected mice.⁴⁸ Finally, it is noteworthy that *in vivo* inhibition of cathepsin B improves memory and synaptic transmission in transgenic mice over-expressing APP, interfering with amyloidogenic APP processing.⁴⁹ On the contrary, calpain inhibition is also protective *in vivo* against cognitive loss in another AD animal model-APP/PS1 mice- by upregulating the phosphorylation levels of the transcription factor CREB (cAMP Responsive Element Binding Protein) without any significant change in A β peptide levels.⁵⁰

Caspases and tau. Studies from cellular and animal models indicate that caspases have also been implicated in mechanisms of tau-mediated neurodegeneration in AD.^{51,52} According to this hypothesis, A β peptide promotes neuronal pathological tau filament assembly by triggering caspases activation leading to tau cleavage.⁵³ This event, in turn, generates a proteolytic products that assemble more rapidly and extensively into tau pathological filaments.^{54,55} Aberrant activation of caspase(s), following apoptotic stimuli or neurodegenerative insults, may produce one or more toxic NH₂-tau fragments, that further contribute to propagate and increase cellular dysfunctions in AD.^{56,57} Colocalization of hyperphosphorylated tau and active caspase-3 and 6 has been recently detected in brainstem of young and old AD patients.²⁶ The finding that the rTg4510 tau transgenic mouse shows caspase-3 activation provides additional supporting evidence linking caspase-3 and tau-mediated neurodegeneration.⁵⁸ Caspase-9 activation and caspase-cleaved tau forms have been documented in AD hippocampal brain sections.¹⁸ Finally over-expression of Bcl-2 in a triple transgenic Alzheimer mouse model harboring PS1(M146V), APP(Swe) and tau(P301L) transgenes limits caspase-9 activation, attenuates the processing of APP and tau thus reducing the number of NFTs and extracellular deposits of A β associated with the progression of this disease.⁵⁹

It remains to be determined if frank apoptosis is a necessary and early event in the neurodegeneration. According to this view, a positive feedback loop in neurodegeneration would be activated whereby caspase(s) generate A β , which in turn exerts a noxious action on tau proteins and further activates caspase(s) in neighboring neurons eventually dying by apoptosis. In this context, other modes of cell death could contribute to neuronal loss in AD⁶⁰ and other proteases, such as calpain and cathepsin, can be also directly or indirectly activated by caspases during apoptosis.⁶¹ Finally, an intricate cross-talk between these proteases systems has been reported during apoptosis of neuronal cells.⁶² Thus, although other caspase-independent pathways may contribute to the AD progression, the *in vivo* treatment with specific caspase(s) inhibitors, which are able to penetrate the blood-brain barrier, may still offer an useful and alternative therapeutic strategy to delay selective neuronal loss associated to such neurodegenerative disease.

The Cerebellar Granule Cells (CGC) Model

A decade ago, our research group hypothesized a possible tight link between improper activation of apoptosis and events related to AD. Cerebellar Granule Neurons (CGNs) from 8 days old rat require depolarizing potassium concentration (25 mM K⁺) for an optimal survival, when explanted in vitro. Upon reduction of extracellular potassium concentration to a more physiological concentration of 5 mM, these neurons progressively undergo apoptosis⁶³ which is largely blocked by neuroprotective agents able to increase calcium influx.⁶⁴ It has been hypothesized that in vitro depolarizing conditions are necessary to maintain intracellular high levels of free calcium, thus mimicking the in vivo situation of continuous electrical stimulation related to the developmental establishment of excitatory synapses originating from mossy fibers.^{9,65} The apoptotic process, as well as nuclear and mitochondrial damage, are reversible up to 4–8 hours of induction suggesting that no rescue is possible even if CGNs are returned to high K⁺ medium.^{66,67} Activation of caspase-3 has been reported after serum/K⁺ starvation⁶⁸ and cell death is attenuated by the selective caspase-3 inhibitor z-DEVD-fmk;^{69,70} although the main effect of such caspase is on DNA fragmentation and chromatin condensation rather than preventing apoptosis.⁷¹ Such conflicting data may reflect the finding that neuronal apoptosis triggered by potassium reduction involves a more intricate caspase(s) activation cascade⁷² and a cross-talk between caspase(s) and other protease(s) further complicates the death signaling.^{73–76} Neurotrophin and physiological neuropeptides, such as IGF, bFGF, BDNF, PACAP, SP and cAMP^{63,64,76–79} also exert their protective action in this neuronal paradigm through different mechanism including the activation of PI3-kinase/Akt pathway;^{76,78,80,81} the stimulation of PKA and/or MAP kinases signaling.^{82,83}

We have been reported that the pro-apoptotic shifting to a low potassium medium activates an amyloidogenic process, which rapidly and irreversibly leads to an unbalance between the “physiological” α -secretase-mediated pathway and the β - α -secretase mediated increased production of A β .⁸⁴ Moreover, the monomeric and oligomeric forms of 4-kDa A β are significantly higher in depolarization-stimulated secretion compared with controls. Such increments are paralleled by a corresponding increase of the β -APPs/ α -APPs ratio in apoptotic secretion, without any significant change of intracellular full-length APP levels. An interesting aspect of such a process is that the released pool of A β may activate a toxic loop that further accelerates and propagates the process of neurodegeneration, affecting neighboring healthy neurons. Indeed, co-incubation of apoptotic cultures with antibodies directed against A β significantly slows down the extension of cell death and quantitatively increases the neuronal survival rate by approximately 50%,^{85,86} thus suggesting that A β peptides may act as soluble and diffusible apoptotic death mediators.

Contextually to the significant increase of amyloidogenic metabolism of APP, also tau undergoes post-translational modifications. After 6 h of potassium deprivation, a change in tau phosphorylation state and caspase(s) and calpain-mediated cleavage occurs in concomitance with a progressive disassembly of cytoskeleton, eventually leading to the generation of a 17 kDa fragment which accumulates in the perikarya of dying cells.⁷³ Furthermore, following the apoptotic trigger, a reactive oxygen species (ROS) production

and progressively reduced mitochondrial function also contribute to neuronal damage.^{87,88} Superoxide dismutase, N-acetyl-L-cysteine and other free radical scavengers partially protect CGNs from death, improving mitochondrial energy metabolism.^{89,90}

The bulk of studies on CGNs, apoptosis and events related to AD prospected a first, consistent positive answer to their possible link. Nevertheless, the observation that these neurons are not the most vulnerable population affected in AD and that few clinical signs of cerebellar anatomopathological dysfunction have been reported in AD patients leaves room for some criticisms about its fully usefulness as in vitro model for this human neurodegenerative disease.

NGF-Deprived PC12 and Hippocampal/Cortical Neuronal Models

NGF (Nerve Growth Factor) is the first neurotrophin to be discovered and is not only endowed with the property of inducing growth of nerve fibers in target neurons, but also of supporting their life via its antiapoptotic action.⁹¹ Numerous in vitro and in vivo data suggest a tight causal relationship between an imbalance in NGF receptor signaling, the activation of amyloidogenic pathway and altered tau metabolism in onset and progression of AD-like neurodegeneration.

TrkA, the high affinity NGF receptor, has been found decreased in the basal forebrain^{92–97} and in the cortex.^{98–100} A switch from TrkA to p75, the low affinity death receptor, it has been described during neuronal aging resulting in increased amyloidogenic processing of APP.^{101–103} p75NTR expression has been directly linked to changes occurring in AD,¹⁰⁴ including the death of basal forebrain neurons,^{105,106} hypothesized to occur through a direct binding of oligomeric A β 1–42 peptides to p75.^{103,107,108} Moreover, some evidences have previously showing a transcriptional p75-mediated regulation on the APP promoter leading to an increase of secreted amyloid precursor protein (sAPP)^{109,110} in neurons.

Several studies report a regulative role of NGF on tau phosphorylation. Stimulation of undifferentiated PC-12 with NGF causes a dephosphorylation of tau proteins,¹¹¹ although an increase of Gsk3 β -mediated tau phosphorylation has also been observed. Interestingly, this tau phosphorylation at defined sites might be required for proper anterograde organelle/mitochondrial transport in differentiated cells.¹¹² On the other hand, NGF deprivation in differentiated PC12 induces apoptosis and hyperphosphorylation both of tau and membrane-bound high molecular weight (HMW) tau, especially in the neuritis. These changes are accompanied by an impairment of its microtubule binding ability and a marked decrease of its solubility. However, in the last stages of apoptosis, tau is dephosphorylated in dying neuronal PC12.^{113,114} In addition, in this apoptotic neuronal model, NGF deprivation also causes an early, caspase-mediated tau cleavage at NH₂ domain with the appearance of the 20–22 kDa tau fragment¹¹⁵ which has been previously demonstrated to be markedly neurotoxic in vitro when overexpressed in primary neuronal cultures.¹¹⁶ NGF might control the endogenous tau protein levels, regulating its metabolism via proteasomal degradation, as demonstrated by NGF-dependent ubiquitination of tau in cultured cells.¹¹⁷ Finally, several evidences support the hypothesis that the role of tau in axonal transport might affect NGF-TrkA signaling in vivo. Indeed, experimental data from retrograde labeling of basal forebrain neurons after injection of fluorogold into multiple sites in cortex and

hippocampus, report that an altered compartmentalization of phosphatase, GSK3 and TrkA immunoreactivity may be responsible for the failure of axonal trafficking and lack of trophic support in aged cholinergic cells.^{118,119}

The hypothesis that a chronic NGF deprivation may be one of the factors involved in the etiology of sporadic forms of AD is validated by the findings that acute treatment with NGF or acetylcholine esterase (AChE) inhibitors, such as ganstigmine and donepezil, rescues the cholinergic and behavioral deficit in AD11 mice. These mice are an in vivo AD transgenic model, in which the phenotypic knockout of NGF is achieved by the expression of recombinant neutralizing antibodies.^{120,121} Finally, clinical encouraging data from ongoing gene therapy trial using NGF-grafted autologous fibroblasts injected into the basal nucleus of Meynert (nbM),¹²² further validate the rationale for the therapeutic administration of human recombinant NGF in AD patients.¹²³

In view of these findings, we carried out a set of experiments in NGF-deprived differentiated PC12 cells^{124,115} and described the crucial steps linking NGF withdrawal, activation of amyloidogenesis, tau truncation and caspase(s)-mediated execution of neuronal death. These studies have been replicated in primary hippocampal and cortical neurons showing that, upon NGF removal, the amyloidogenic pathway is activated with consequent intra and extracellular accumulation of A β peptides and apoptotic death. The overproduced A β is partly released in the culture medium, where it aggregates to form structures largely reminiscent of those forming senile plaques, and in part aggregates within neurons. All these events are prevented by β and α secretase inhibitors, by antibodies directed against A β peptides, or by partial silencing of APP mRNA, whereas they are mimicked by A β 1–42 peptide exposure. Conversely, neurons deprived of serum largely die but, although the amyloidogenic pathway is activated, the exposure to anti A β antibodies does not protect from apoptotic death, further suggesting that the activation of amyloidogenesis following NGF withdrawal is not a simple consequence of an apoptotic trigger but it is strictly related to lack of NGF supply.¹²⁵

In the same experimental model we have also demonstrated an early involvement of tau protein which, under NGF deprivation, undergoes GSK3 β mediated hyperphosphorylation at pathogenic amino acids such as Ser 262 and Thr 231, and is subsequently degraded generating a toxic NH-2-derived 220 amino acid peptide.¹¹⁶ Such tau hyperphosphorylation, as well as apoptotic death, is blocked by A β antibodies or by specific β and/or α -secretases inhibitors and is mimicked by A β 1–42 peptide, suggesting that A β species are the initial trigger. Tau subsequently detaches from microtubules, thus shifting the equilibrium toward its disassembled state and indirectly affecting the whole axonal transport, eventually leading to apoptotic death (Amadoro et al. submitted). Once tau is displaced from microtubules, it would be further phosphorylated at other fibrillogenic site and/or cleaved by proteases (i.e., caspase(s) and calpain), causing disruption of microtubule transport along axons and consequent synaptic dysfunction. All these events are summarized in the Figure 1.

To our knowledge, the NGF-deprived hippocampal culture is presently the only in vitro model whereby both APP, tau altered processing and apoptosis, have been investigated together under strictly controlled conditions. Regarding the direct role of caspases




Figure 1. Schematic representation of the apoptotic mechanisms by which the interruption of NGF signaling affects APP processing and Tau metabolism in hippocampal neurons. For more details see text.

in neuronal in vitro models whose viability is strictly dependent on NGF, an involvement of such proteases in apoptotic cell death caused by NGF deprivation^{126,127} and in the p75-mediated cell death caused by exogenous addition of A β to neuroblastoma cells¹²⁸ has been largely documented. The studies performed in NGF-deprived PC12 cells show that, among all caspase(s) inhibitors tested, only blockage of executor caspases 2, 12, 6 and 8 exerts an almost total protection from death and from A β production, whereas inhibition of effector caspase 3 does not exert a similar action.¹²⁴ In a similar fashion and in agreement with others,^{129,130} treatment with Z-DEVD-fmk, a specific inhibitor of caspase-3 only partially rescued hippocampal neurons from death, probably because this protease is not activated at early times upon NGF withdrawal in this neuronal paradigm (Amadoro et al. submitted). On the contrary, pharmacological inhibition of caspase 3 markedly inhibited caspases-mediated tau cleavage, without any significant effects on GSK3 β -mediated tau hyperphosphorylation (Amadoro et al. submitted). Moreover, the finding that the general cell-permeable caspase inhibitor z-VAD does not significantly affect ThT-positive A β structures production in NGF-deprived PC12, whereas partially rescues cells from apoptotic death,¹²⁴ delineates a complex chain of events between NGF withdrawal, A β production, apoptosis and tau modifications. As mentioned above, the causal and temporal relationship between caspases-mediated cell death and APP processing appears cell-specific and signaling-dependent and probably initiates a toxic cycle of cellular A β production/neuronal loss, which is difficult to elucidate in its actual sequence. Thus, although elevated A β may lead to apoptotic cell death after injury or disease and caspase(s) inhibition may protect against this event, a causal relationship could not be proven as blockade of caspase(s) might also prevent tau modifications and cell death unrelated to A β toxicity. Further investigations aimed at selectively reducing A β levels, without targeting caspase(s) activity (i.e., by directly altering α , β and/or α secretase activity), will provide additional insights into this cascade to definitively establish if apoptosis is the primary cause of A β production/tau modification or is it a sort of downstream consequence, eventually ending in cell death.

Acknowledgements

Financial support was received from the Regione Lazio, FIRB 2003, Prin 2006 and INRCA 2008 (Istituto Nazionale Riposo e Cura per Anziani) to P. Callisano.

References

- Selkoe DJ. Alzheimer's disease: genes, proteins and therapy. *Physiol Rev* 2001; 81:741-66.
- Shen Y, He P, Zhong Z, McAllister C, Lindholm K. Distinct destructive signal pathways of neuronal death in Alzheimer's disease. *Trends Mol Med* 2006; 12:574-9.
- Salehi A, Delcroix JD, Mobley WC. Traffic at the intersection of neurotrophic factor signaling and neurodegeneration. *Trends Neurosci* 2003; 26:73-80.
- Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 2002; 297:353-6.
- Blurton-Jones M, Laferla FM. Pathways by which Abeta facilitates tau pathology. *Curr Alzheimer Res* 2006; 3:437-48.
- Mattson MP. Apoptosis in neurodegenerative disorders. *Nat Rev Mol Cell Biol* 2000; 1:120-9.
- Mattson MP, Magnus T. Ageing and neuronal vulnerability. *Nat Rev Neurosci* 2006; 7:278-94.
- Schulte-Herbrüggen O, Braun A, Rochlitz S, Jockers-Scherübl MC, Hellweg R. Neurotrophic factors—a tool for therapeutic strategies in neurological, neuropsychiatric and neuroimmunological diseases? *Curr Med Chem* 2007; 14:2318-29.
- Canu N, Callisano P. In vitro cultured neurons for molecular studies correlating apoptosis with events related to Alzheimer disease. *Cerebellum* 2003; 2:270-8.
- Shimohama S. Apoptosis in Alzheimer's disease—an update. *Apoptosis* 2000; 5:9-16.
- Behl C. Apoptosis and Alzheimer's disease. *J Neural Transm* 2000; 107:1325-44.
- Kitamura Y, Shimohama S, Kamoshima W, Ota T, Matsuoka Y, Nomura Y, et al. Alteration of proteins regulating apoptosis, Bcl-2, Bcl-x, Bax, Bak, Bad, ICH-1 and CPP32, in Alzheimer's disease. *Brain Res* 1998; 780:260-9.
- Su JH, Deng G, Cotman CW. Bax protein expression is increased in Alzheimer's brain: correlations with DNA damage, Bcl-2 expression and brain pathology. *J Neuropathol Exp Neurol* 1997; 56:86-93.
- Stadelmann C, Deckwerth TL, Srinivasan A, Bancher C, Brück W, Jellinger K, et al. Activation of caspase-3 in single neurons and autophagic granules of granulovacuolar degeneration in Alzheimer's disease. Evidence for apoptotic cell death. *Am J Pathol* 1999; 155:1459-66.
- Selznick LA, Holtzman DM, Han BH, Gökden M, Srinivasan AN, Johnson EM Jr, et al. In situ immunodetection of neuronal caspase-3 activation in Alzheimer disease. *J Neuropathol Exp Neurol* 1999; 58:1020-6.
- Guo H, Albrecht S, Bourdeau M, Petzke T, Bergeron C, LeBlanc AC. Active caspase-6 and caspase-6-cleaved tau in neuropil threads, neuritic plaques and neurofibrillary tangles of Alzheimer's disease. *Am J Pathol* 2004; 165:523-31.
- Rohn TT, Head E, Nesse WH, Cotman CW, Cribbs DH. Activation of caspase-8 in the Alzheimer's disease brain. *Neurobiol Dis* 2001; 8:1006-16.
- Rohn TT, Rissman RA, Davis MC, Kim YE, Cotman CW, Head E. Caspase-9 activation and caspase cleavage of tau in the Alzheimer's disease brain. *Caspase-9 activation and caspase cleavage of tau in the Alzheimer's disease brain. Neurobiol Dis* 2002; 11:341-54.
- Wu CK, Thal L, Pizzo D, Hansen L, Masliah E, Geula C. Apoptotic signals within the basal forebrain cholinergic neurons in Alzheimer's disease. *Exp Neurol* 2005; 195:484-96.
- Albrecht S, Bourdeau M, Bennett D, Mufson EJ, Bhattacharjee M, LeBlanc AC. Activation of caspase-6 in aging and mild cognitive impairment. *Am J Pathol* 2007; 170:1200-9.
- Suzuki T, Nishiyama K, Yamamoto A, Inazawa J, Iwaki T, Yamada T, et al. Molecular cloning of a novel apoptosis-related gene, human Nap1 (NCKAP1), and its possible relation to Alzheimer disease. *Genomics* 2000; 63:246-54.
- Cribbs DH, Poon WW, Rissman RA, Blurton-Jones M. Caspase-mediated degeneration in Alzheimer's disease. *Am J Pathol* 2004; 165:353-5.
- Gastard MC, Troncoso JC, Koliatsos VE. Caspase activation in the limbic cortex of subjects with early Alzheimer's disease. *Ann Neurol* 2003; 54:393-8.
- Louneva N, Cohen JW, Han LY, Talbot K, Wilson RS, Bennett DA, et al. Caspase 3 is enriched in postsynaptic densities and increased in Alzheimer's disease. *Am J Pathol* 2008; 173:1488-95.
- Su JH, Kesslak JP, Head E, Cotman CW. Caspase-cleaved amyloid precursor protein and activated caspase-3 are co-localized in the granules of granulovacuolar degeneration in Alzheimer's disease and Down's syndrome brain. *Acta Neuropathol* 2002; 104:1-6.
- Wai MS, Liang Y, Shi C, Cho EY, Kung HF, Yew DT. Co-localization of hyperphosphorylated tau and caspases in the brainstem of Alzheimer's disease patients. *Biogerontology* 2008; In press.
- Gervais FG, Xu D, Robertson GS, Vaillancourt JP, Zhu Y, Huang J, et al. Involvement of caspases in proteolytic cleavage of Alzheimer's amyloid-beta precursor protein and amyloidogenic A beta peptide formation. *Cell* 1999; 97:395-406.
- Lu DC, Rabizadeh S, Chandra S, Shayya RF, Ellerby LM, Ye X, et al. A second cytotoxic proteolytic peptide derived from amyloid beta-protein precursor. *Nat Med* 2000; 6:397-404.
- Tesco G, Koh YH, Kang EL, Cameron AN, Das S, Sena-Estevés M, et al. Depletion of GGA3 stabilizes BACE and enhances beta-secretase activity. *Neuron* 2007; 54:721-37.
- Marín N, Romero B, Bosch-Morell F, Llansola M, Felipe V, Romá J, et al. Beta-amyloid-induced activation of caspase-3 in primary cultures of rat neurons. *Mech Ageing Dev* 2000; 119:63-7.
- Loo DT, Copani A, Pike CJ, Whittemore ER, Walencewicz AJ, Cotman CW. Apoptosis is induced by beta-amyloid in cultured nervous system neurons. *Proc Natl Acad Sci USA* 1993; 90:7951-5.
- Uetsuki T, Takemoto K, Nishimura I, Okamoto M, Niinobe M, Momoi T, et al. Activation of neuronal caspase-3 by intracellular accumulation of wild-type Alzheimer amyloid precursor protein. *J Neurosci* 1999; 19:6955-64.
- Nishimura I, Uetsuki T, Kuwako K, Hara T, Kawakami T, Aimoto S, et al. Cell death induced by a caspase-cleaved transmembrane fragment of the Alzheimer amyloid precursor protein. *Cell Death Differ* 2002; 9:199-208.
- Chan SL, Mattson MP. Caspase and calpain substrates: roles in synaptic plasticity and cell death. *J Neurosci Res* 1999; 58:167-90.
- Glabe C. Intracellular mechanisms of amyloid accumulation and pathogenesis in Alzheimer's disease. *J Mol Neurosci* 2001; 17:137-45.
- Oddo S, Caccamo A, Shepherd JD, Murphy MP, Golde TE, Kaye R, et al. Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular Abeta and synaptic dysfunction. *Neuron* 2003; 39:409-21.
- Lustbader JW, Cirilli M, Lin C, Xu HW, Takuma K, Wang N, et al. ABAD directly links Abeta to mitochondrial toxicity in Alzheimer's disease. *Science* 2004; 304:448-52.
- Du H, Guo L, Fang F, Chen D, Sosunov AA, McKhann GM, et al. Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med* 2008; 14:1097-105.
- Oddo S. The ubiquitin-proteasome system in Alzheimer's disease. *J Cell Mol Med* 2008; 12:363-73.
- Song S, Lee H, Kam TI, Tai ML, Lee JY, Noh JY, et al. E2-25K/Hip-2 regulates caspase-12 in ER stress-mediated Abeta neurotoxicity. *J Cell Biol* 2008; 82:675-84.
- Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, et al. Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 2000; 403:98-103.
- Ivins KJ, Thornton PL, Rohn TT, Cotman CW. Neuronal apoptosis induced by beta-amyloid is mediated by caspase-8. *Neurobiol Dis* 1999; 6:440-9.
- Giovanni A, Keramaris E, Morris EJ, Hou ST, O'Hare M, Dyson N, et al. E2F1 mediates death of B-amyloid-treated cortical neurons in a manner independent of p53 and dependent on Bax and caspase 3. *J Biol Chem* 2000; 275:11553-60.
- Troy CM, Rabacchi SA, Friedman WJ, Frappier TF, Brown K, Shelanski ML. Caspase-2 mediates neuronal cell death induced by beta-amyloid. *J Neurosci* 2000; 20:1386-92.
- Wei Z, Song MS, MacTavish D, Jhamandas JH, Kar S. Role of calpain and caspase in beta-amyloid-induced cell death in rat primary septal cultured neurons. *Neuropharmacology* 2008; 54:721-33.
- LeBlanc A, Liu H, Goodyer C, Bergeron C, Hammond J. Caspase-6 role in apoptosis of human neurons, amyloidogenesis and Alzheimer's disease. *J Biol Chem* 1999; 274:23426-36.
- Weidemann A, Paliga K, Dürrwang U, Reinhard FB, Schuckert O, Evin G, et al. Proteolytic processing of the Alzheimer's disease amyloid precursor protein within its cytoplasmic domain by caspase-like proteases. *J Biol Chem* 1999; 274:5823-9.
- Abrahamson EE, Ikonomic MD, Ciallella JR, Hope CE, Paljug WR, Isanski BA, et al. Caspase inhibition therapy abolishes brain trauma-induced increases in Abeta peptide: implications for clinical outcome. *Exp Neurol* 2006; 197:437-50.
- Hook VY, Kindy M, Hook G. Inhibitors of cathepsin B improve memory and reduce beta-amyloid in transgenic Alzheimer disease mice expressing the wild-type, but not the Swedish mutant, beta-secretase site of the amyloid precursor protein. *J Biol Chem* 2008; 283:7745-53.
- Trinchese F, Fa' M, Liu S, Zhang H, Hidalgo A, Schmidt SD, et al. Inhibition of calpains improves memory and synaptic transmission in a mouse model of Alzheimer disease. *J Clin Invest* 2008; 118:776-807.
- Dickson DW. Apoptotic mechanisms in Alzheimer neurofibrillary degeneration: cause or effect? *J Clin Invest* 2004; 114:23-7.
- García-Sierra F, Mondragón-Rodríguez S, Basurto-Islas G. Truncation of tau protein and its pathological significance in Alzheimer's disease. *J Alzheimers Dis* 2008; 14:401-9.
- Cotman CW, Poon WW, Rissman RA, Blurton-Jones M. The role of caspase cleavage of tau in Alzheimer disease neuropathology. *J Neuropathol Exp Neurol* 2005; 64:104-12.
- Gamblin TC, Chen F, Zambrano A, Abrahá A, Lagalwar S, Guillozet AL, et al. Caspase cleavage of tau: linking amyloid and neurofibrillary tangles in Alzheimer's disease. *Proc Natl Acad Sci USA* 2003; 100:10032-7.
- Cho JH, Johnson GV. Glycogen synthase kinase 3 beta induces caspase-cleaved tau aggregation in situ. *J Biol Chem* 2004; 279:54716-23.
- Fasulo L, Ugolini G, Visintin M, Bradbury A, Brancolini C, Verzillo V, et al. The neuronal microtubule-associated protein tau is a substrate for caspase-3 and an effector of apoptosis. *J Neurochem* 2000; 75:624-33.
- Chung CW, Song YH, Kim IK, Yoon WJ, Ryu BR, Jo DG, et al. Proapoptotic effects of tau cleavage product generated by caspase-3. *Neurobiol Dis* 2001; 8:162-72.
- Ramálho RM, Viana RJ, Castro RE, Steer CJ, Low WC, Rodríguez CM. Apoptosis in transgenic mice expressing the P301L mutated form of human tau. *Mol Med* 2008; 1485:309-17.
- Rohn TT, Vyas V, Hernandez-estrada T, Nichol KE, Christie LA, Head E. Lack of pathology in a triple transgenic mouse model of Alzheimer's disease after overexpression of the anti-apoptotic protein bcl-2. *J Neurosci* 2008; 28:3051-9.

60. Zilkova M, Koson P, Zilka N. The hunt for dying neurons: insight into the neuronal loss in Alzheimer's disease. *Bratisl Lek Listy* 2006; 107:366-73.
61. Yuan J. Divergence from a dedicated cellular suicide mechanism: exploring the evolution of cell death. *Mol Cell* 2006; 23:1-12.
62. Bredesen DE, Rao RV, Mehlen P. Cell death in the nervous system. *Nature* 2006; 443:796-802.
63. D'Mello SR, Galli C, Ciotti T, Calissano P. Induction of apoptosis in cerebellar granule neurons by low potassium: inhibition of death by insulin-like growth factor I and cAMP. *Proc Natl Acad Sci USA* 1993; 90:10989-93.
64. Galli C, Meucci O, Scorziello A, Werge TM, Calissano P, Schettini G. Apoptosis in cerebellar granule cells is blocked by high KCl, forskolin and IGF-1 through distinct mechanisms of action: the involvement of intracellular calcium and RNA synthesis. *J Neurosci* 1995; 15:1172-9.
65. Borsello T, Di Luzio A, Ciotti MT, Calissano P, Galli C. Granule neuron DNA damage following deafferentation in adult rats cerebellar cortex: a lesion model. *Neuroscience* 2000; 95:163-71.
66. Nardi N, Avidan G, Daily D, Zilkha-Falb R, Barzilai A. Biochemical and temporal analysis of events associated with apoptosis induced by lowering the extracellular potassium concentration in mouse cerebellar granule neurons. *J Neurochem* 1997; 68:750-9.
67. Schulz JB, Beinroth S, Weller M, Wüllner U, Klockgether T. Endonucleolytic DNA fragmentation is not required for apoptosis of cultured rat cerebellar granule neurons. *Neurosci Lett* 1998; 245:9-12.
68. Armstrong RC, Aja TJ, Hoang KD, Gaur S, Bai X, Alnemri ES, Litwack G, Karanewsky DS, Fritz LC, Tomaselli KJ. Activation of the CED3/ICE-related protease CPP32 in cerebellar granule neurons undergoing apoptosis but not necrosis. *J Neurosci* 1997; 17:553-62.
69. Eldadah BA, Yakovlev AG, Faden AI. The role of CED-3-related cysteine proteases in apoptosis of cerebellar granule cells. *J Neurosci* 1997; 17:6105-13.
70. Eldadah BA, Ren RF, Faden AI. Ribozyme-mediated inhibition of caspase-3 protects cerebellar granule cells from apoptosis induced by serum-potassium deprivation. *J Neurosci* 2000; 20:179-86.
71. D'Mello SR, Kuan CY, Flavell RA, Rakic P. Caspase-3 is required for apoptosis-associated DNA fragmentation but not for cell death in neurons deprived of potassium. *J Neurosci Res* 2000; 59:24-31.
72. Gerhardt E, Kügler S, Leist M, Beier C, Berliocchi L, Volbracht C, et al. Cascade of caspase activation in potassium-deprived cerebellar granule neurons: targets for treatment with peptide and protein inhibitors of apoptosis. *Mol Cell Neurosci* 2001; 4:717-31.
73. Canu N, Dus L, Barbato C, Ciotti MT, Brancolini C, Rinaldi AM, et al. Tau cleavage and dephosphorylation in cerebellar granule neurons undergoing apoptosis. *J Neurosci* 1998; 18:7061-74.
74. Canu N, Tufi R, Serafino AL, Amadoro G, Ciotti MT, Calissano P. Role of the autophagolysosomal system on low potassium-induced apoptosis in cultured cerebellar granule cells. *J Neurochem* 2005; 92:1228-42.
75. Kaasik A, Rikk T, Piirsoo A, Zharkovsky T, Zharkovsky A. Upregulation of lysosomal cathepsin L and autophagy during neuronal death induced by reduced serum and potassium. *Eur J Neurosci* 2005; 22:1023-31.
76. Amadoro G, Pieri M, Ciotti MT, Carunchio I, Canu N, Calissano P, et al. Substance P provides neuroprotection in cerebellar granule cells through Akt and MAPK/Erk activation: evidence for the involvement of the delayed rectifier potassium current. *Neuropharmacology* 2007; 52:1366-77.
77. Kubo T, Nonomura T, Enokido Y, Hatanaka H. Brain-derived neurotrophic factor (BDNF) can prevent apoptosis of rat cerebellar granule neurons in culture. *Brain Res Dev Brain Res* 1995; 85:249-58.
78. D'Mello SR, Borodetz K, Soltoff SP. Insulin-like growth factor and potassium depolarization maintain neuronal survival by distinct pathways: possible involvement of PI 3-kinase in IGF-1 signaling. *J Neurosci* 1997; 17:1548-60.
79. Journot L, Villalba M, Bockaert J. PACAP-38 protects cerebellar granule cells from apoptosis. *Ann NY Acad Sci* 1998; 865:100-10.
80. Dudek H, Datta SR, Franke TF, Birnbaum MJ, Yao R, Cooper GM, et al. Regulation of neuronal survival by the serine-threonine protein kinase Akt. *Science* 1997; 275:661-5.
81. Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, et al. Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 1999; 96:857-68.
82. Kienlen Campard P, Crochemore C, René F, Monnier D, Koch B, Loeffler JP. DNA PACAP type I receptor activation promotes cerebellar neuron survival through the cAMP/PKA signaling pathway. *Cell Biol* 1997; 16:323-33.
83. Villalba M, Bockaert J, Journot L. Pituitary adenylate cyclase-activating polypeptide (PACAP-38) protects cerebellar granule neurons from apoptosis by activating the mitogen-activated protein kinase (MAP kinase) pathway. *J Neurosci* 1997; 17:83-90.
84. Galli C, Piccini A, Ciotti MT, Castellani L, Calissano P, Zaccheo D, et al. Increased amyloidogenic secretion in cerebellar granule cells undergoing apoptosis. *Proc Natl Acad Sci USA* 1998; 95:1247-52.
85. Piccini A, Ciotti MT, Vitolo OV, Calissano P, Tabaton M, Galli C. Endogenous APP derivatives oppositely modulate apoptosis through an autocrine loop. *Neuroreport* 2000; 11:1375-9.
86. De Berardinis M, Ciotti MT, Amadoro G, Galli C, Calissano P. Transfer of the apoptotic message in sister cultures of cerebellar neurons. *Neuroreport* 2001; 12:2137-40.
87. Atlante A, Gagliardi S, Marra E, Calissano P. Neuronal apoptosis in rats is accompanied by rapid impairment of cellular respiration and is prevented by scavengers of reactive oxygen species. *Neurosci Lett* 1998; 245:127-30.
88. Atlante A, Bobba A, Calissano P, Passarella S, Marra E. The apoptosis/necrosis transition in cerebellar granule cells depends on the mutual relationship of the antioxidant and the proteolytic systems which regulate ROS production and cytochrome *c* release en route to death. *J Neurochem* 2003; 84:960-71.
89. Atlante A, Passarella S. Detection of reactive oxygen species in primary cultures of cerebellar granule cells. *Brain Res Brain Res Protoc* 1999; 4:266-70.
90. Schulz JB, Weller M, Klockgether T. Potassium deprivation-induced apoptosis of cerebellar granule neurons: a sequential requirement for new mRNA and protein synthesis, ICE-like protease activity, and reactive oxygen species. *J Neurosci* 1996; 16:4696-706.
91. Levi-Montalcini R. Nerve growth factor. *Science* 1975; 187:113.
92. Boissière F, Hunot S, Faucheux B, Hersh LB, Agid Y, Hirsch EC. Trk neurotrophin receptors in cholinergic neurons of patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 1997; 8:1-8.
93. Chu Y, Cochran EJ, Bennett DA, Mufson EJ, Kordower JH. Downregulation of trkA mRNA within nucleus basalis neurons in individuals with mild cognitive impairment and Alzheimer's disease. *J Comp Neurol* 2001; 437:296-307.
94. Ginsberg SD, Che S, Wu J, Counts SE, Mufson EJ. Downregulation of trk but not p75NTR gene expression in single cholinergic basal forebrain neurons mark the progression of Alzheimer's disease. *J Neurochem* 2006; 97:475-87.
95. Mufson EJ, Lavine N, Jaffar S, Kordower JH, Quirion R, Saragovi HU. Reduction in p140-TrkA receptor protein within the nucleus basalis and cortex in Alzheimer's disease. *Exp Neurol* 1997; 146:91-103.
96. Mufson EJ, Ma SY, Cochran EJ, Bennett DA, Beckett LA, Jaffar S, et al. Loss of nucleus basalis neurons containing trkA immunoreactivity in individuals with mild cognitive impairment and early Alzheimer's disease. *J Comp Neurol* 2000; 427:19-30.
97. Salehi A, Verhaagen J, Dijkhuizen PA, Swaab DF. Co-localization of high-affinity neurotrophin receptors in nucleus basalis of Meynert neurons and their differential reduction in Alzheimer's disease. *Neuroscience* 1996; 75:373-87.
98. Counts SE, Nadeem M, Wu J, Ginsberg SD, Saragovi HU, Mufson EJ. Reduction of cortical TrkA but not p75(NTR) protein in early-stage Alzheimer's disease. *Ann Neurol* 2004; 56:520-31.
99. Hock C, Heese K, Hulette C, Rosenberg C, Otten U. Region-specific neurotrophin imbalances in Alzheimer disease: decreased levels of brain-derived neurotrophic factor and increased levels of nerve growth factor in hippocampus and cortical areas. *Arch Neurol* 2000; 57:846-51.
100. Savaskan E, Müller-Spahn F, Olivieri G, Bruttel S, Otten U, Rosenberg C, et al. Alterations in trkA, trkB and trkC receptor immunoreactivities in parietal cortex and cerebellum in Alzheimer's disease. *Eur Neurol* 2000; 44:172-80.
101. Costantini C, Weindrich R, Della Valle G, Puglielli L. A TrkA-to-p75NTR molecular switch activates amyloid beta-peptide generation during aging. *Biochem J* 2005; 391:59-67.
102. Costantini C, Scrabble H, Puglielli L. An aging pathway controls the TrkA to p75NTR receptor switch and amyloid beta-peptide generation. *EMBO J* 2006; 25:1997-2006.
103. Costantini C, Rossi F, Formaggio E, Bernardoni R, Cecconi D, Della-Bianca V. Characterization of the signaling pathway downstream p75 neurotrophin receptor involved in beta-amyloid peptide-dependent cell death. *J Mol Neurosci* 2005; 25:141-56.
104. Schliebs R. Basal forebrain cholinergic dysfunction in Alzheimer's disease—interrelationship with beta-amyloid, inflammation and neurotrophin signaling. *Neurochem Res* 2005; 30:895-908.
105. Yeo TT, Chua-Couzens J, Butcher LL, Bredesen DE, Cooper JD, Valletta JS, et al. Absence of p75NTR causes increased basal forebrain cholinergic neuron size, choline acetyltransferase activity, and target innervation. *J Neurosci* 1997; 17:7594-605.
106. Volosin M, Song W, Almeida RD, Kaplan DR, Hempstead BL, Friedman WJ. Interaction of survival and death signaling in basal forebrain neurons: roles of neurotrophins and proneurotrophins. *J Neurosci* 2006; 26:7756-66.
107. Yaar M, Arble BL, Stewart KB, Qureshi NH, Kowall NW, Gilchrist BA. p75 (NTR) antagonistic cyclic peptide decreases the size of beta amyloid-induced brain inflammation. *Cell Mol Neurobiol* 2008; 28:1027-31.
108. Sotthibundhu A, Sykes AM, Fox B, Underwood CK, Thangnipon W, Coulson EJ. Beta-amyloid (1-42) induces neuronal death through the p75 neurotrophin receptor. *J Neurosci* 2008; 28:3941-6.
109. Ge YW, Lahiri DK. Regulation of promoter activity of the APP gene by cytokines and growth factors: implications in Alzheimer's disease. *Ann NY Acad Sci* 2002; 973:463-7.
110. Rossner S, Ueberham U, Schliebs R, Perez-Polo JR, Bigl V. p75 and TrkA receptor signaling independently regulate amyloid precursor protein mRNA expression, isoform composition and protein secretion in PC12 cells. *J Neurochem* 1998; 71:757-66.
111. Fisher A, Heldman E, Gurwitz D, Haring R, Karton Y, Meshulam H, et al. M1 agonists for the treatment of Alzheimer's disease. Novel properties and clinical update. *Ann NY Acad Sci* 1996; 777:189-96.
112. Tatebayashi Y, Haque N, Tung YC, Iqbal K, Grundke-Iqbal I. Role of tau phosphorylation by glycogen synthase kinase-3beta in the regulation of organelle transport. *J Cell Sci* 2004; 117:1653-63.
113. Nuydens R, Dispersyn G, de Jong M, van den Kieboom G, Borgers M, Geerts H. Aberrant tau phosphorylation and neurite retraction during NGF deprivation in PC12 cells. *Biochem Biophys Res Commun* 1997; 240:687-91.
114. Shelton SB, Johnson GV. Tau and HMW tau phosphorylation and compartmentalization in apoptotic neuronal PC12 cells. *J Neurosci Res* 2001; 66:203-13.

115. Corsetti V, Amadoro G, Gentile A, Capsoni S, Ciotti MT, Cencioni MT, et al. Identification of a caspase-derived N-terminal tau fragment in cellular and animal Alzheimer's disease models. *Mol Cell Neurosci* 2008; 38:381-92.
116. Amadoro G, Ciotti MT, Costanzi M, Cestari V, Calissano P, Canu N. NMDA receptor mediates tau-induced neurotoxicity by calpain and ERK/MAPK activation. *Proc Natl Acad Sci USA* 2006; 103:2892-7.
117. Babu JR, Geetha T, Wooten MW. Sequestosome 1/p62 shuttles polyubiquitinated tau for proteasomal degradation. *J Neurochem* 2005; 94:192-203.
118. Niewiadomska G, Baksalerska-Pazera M, Riedel G. Altered cellular distribution of phospho-tau proteins coincides with impaired retrograde axonal transport in neurons of aged rats. *Ann NY Acad Sci* 2005; 1048:287-95.
119. Niewiadomska G, Baksalerska-Pazera M, Riedel G. Cytoskeletal transport in the aging brain: focus on the cholinergic system. *Rev Neurosci* 2006; 17:581-618.
120. Ruberti F, Capsoni S, Comparini A, Di Daniel E, Franzot J, Gonfloni S, et al. Phenotypic knockout of nerve growth factor in adult transgenic mice reveals severe deficits in basal forebrain cholinergic neurons, cell death in the spleen and skeletal muscle dystrophy. *J Neurosci* 2000; 20:2589-601.
121. Capsoni S, Cattaneo A. On the molecular basis linking nerve growth factor (NGF) to Alzheimer's disease. *Cell Mol Neurobiol* 2006; 26:619-33.
122. Tuszynski MH, Thal L, Pay M, Salmon DP, U HS, Bakay R, et al. A phase 1 clinical trial of nerve growth factor gene therapy for Alzheimer disease. *Nat Med* 2005; 11:551-5.
123. Cattaneo A, Capsoni S, Paoletti F. Towards non invasive nerve growth factor therapies for Alzheimer's disease. *J Alzheimers Dis* 2008; 15:255-83.
124. Matrone C, Di Luzio A, Meli G, D'Aguzzo S, Severini C, Ciotti MT, et al. Activation of the amyloidogenic route by NGF deprivation induces apoptotic death in PC12 cells. *J Alzheimers Dis* 2008; 13:81-96.
125. Matrone C, Ciotti MT, Mercanti D, Marolda R, Calissano P. NGF and BDNF signaling control amyloidogenic route and Ab production in hippocampal neurons. *Proc Natl Acad Sci USA* 2008; 105:13138-43.
126. Park DS, Morris EJ, Stefanis L, Troy CM, Shelanski ML, Geller HM, et al. Multiple pathways of neuronal death induced by DNA-damaging agents, NGF deprivation and oxidative stress. *J Neurosci* 1998; 18:830-40.
127. Troy CM, Rabacchi SA, Hohl JB, Angelastro JM, Greene LA, Shelanski ML. Death in the balance: alternative participation of the caspase-2 and -9 pathways in neuronal death induced by nerve growth factor deprivation. *J Neurosci* 2001; 21:5007-16.
128. Perini G, Della-Bianca V, Politi V, Della Valle G, Dal-Pra I, et al. Role of p75 neurotrophin receptor in the neurotoxicity by beta-amyloid peptides and synergistic effect of inflammatory cytokines. *J Exp Med* 2002; 195:907-18. Erratum in: *J Exp Med* 2002; 195:1231.
129. Rametti A, Esclaire F, Yardin C, Terro F. Linking alterations in tau phosphorylation and cleavage during neuronal apoptosis. *J Biol Chem* 2004; 279:54518-28.
130. Bhat RV, Leonov S, Luthman J, Scott CW, Lee CM. Interactions between GSK3 β and caspase signalling pathways during NGF deprivation induced cell death. *J Alzheimers Dis* 2002; 4:291-30.